



References for Additional Information on Risk Factors for Late Onset Hearing Loss

NOTE: These references were used at the time information was being collected for EHDDI's Risk Factor Information Sheets. Please note that the online references have not been updated recently and some web links may not be accurate.

EXTENDED STAY IN NICU

1. Merck Manual of Diagnosis & Therapy. Section 7. Ear, Nose, & Throat Disorders. Chapter 85. Inner Ear. <http://www.merck.com/mrkshared/mmanual/section7/chapter85/85j.jsp>
2. Van Naarden, K. and Decoufle, P. Relative and Attributable Risks for Moderate to Profound Bilateral Sensorineural Hearing Impairment Associated with Lower Birth Weight in Children 2 to 10 Years Old. *Pediatrics*. 1999;104;905-910.
3. Nicolosi L, Langella L, Krzyzstofiak A, Ticca F. Streptococcus pneumonia meningitis in children. Case records 1985-2003. [article in Italian] *Infez Med*. 2004 Dec;12(4):252-258.
4. Oysu C, Aslan I, Baserer N. Incidence of cochlear involvement in hyperbilirubinemic deafness. *Ann Otol Rhinol Laryngol*. 2002. Nov;111(11):1021-5.
5. Fligor BJ, Neault MW, Mullen CH, Feldman HA, Jones DT. Factors associated with sensorineural hearing loss among survivors of extracorporeal membrane oxygenation therapy. *Pediatrics*. 2005 Jun; 115 (6):1519-28.
6. North Carolina Late Onset Hearing Loss Detection and Intervention, <http://www.ncnewbornhearing.org/LateOnset/Risk%20Factors.htm>
7. DHHS, Agency for Healthcare Research and Quality, July 2000. no. 237. Apgar scores found to be useful in assessing premature newborns, <http://www.ahrq.gov/research/jul00/0700RA5.htm#head4>
8. Usami, S. et al., Sensorineural Hearing Loss Associated with the Mitochondrial Mutations. *Adv. Otorhinolaryngol* 2000; 56:203-211
9. Subcommittee on Hyperbilirubinemia, Clinical Practice Guideline. Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*. 2004 July. 114(1):297-316.
10. Cash, Sheryl. Guideline Offers Direction for Prompt Diagnosis, Treatment of Hyperbilirubinemia. AAP News, American Academy of Pediatrics, July 2004.

FEATURES ASSOCIATED WITH SYNDROMES

1. Gene Reviews, www.genetests.com
2. Online Mendelian Inheritance in Man, <http://www3.ncbi.nlm.nih.gov/entrez/dispmim.cgi?id=214800>

FAMILY HISTORY OF HEARING LOSS

1. Gene Reviews, Deafness and Hereditary Hearing Loss Overview, <http://www.genetests.com/servlet/access?id=8888891&key=yqUmazOwMMLvG&gry=INSERTGRY&fcn=y&fw=YWtM&filename=/glossary/profiles/deafness-overview/index.html>
2. Virginia Early Hearing Detection & Intervention Services (VEHDIS) <http://www.vahealth.org/hearing/>

CRANIOFACIAL ANOMALIES

1. Virginia Early Hearing Detection and Intervention System (VEHDIS), <http://www.vehdis.info/identifying%20risk%20indicators.htm>
2. http://www.drgreene.com/21_1184.html
3. March of Dimes, http://www.marchofdimes.com/professionals/681_1210.asp
4. Francois M, Weiner-Vacher SR, Falala M, Narcy P. Audiological assessment of infants and children with preauricular tags. *Audiology*. 1995. Jan-Feb;34(1):1-5.
5. Zawadzka-Glos L, Bielicka A, Chmielick M. Assessment of hearing in children in postoperative treatment of choanal atresia. 2002. Vol5, http://www.borgis.pl/czytelnia/new_med/2002/02/05.php
6. Baylor College of Medicine Grand Rounds, <http://www.bcm.edu/oto/grand/62493.html>
7. The National Craniofacial Association, <http://www.faces-cranio.org/Disord/Hemi.htm>

MATERNAL ILLNESS

1. Virginia Early Hearing Detection & Intervention Service (VEHDIS), <http://www.vahealth.org/hearing.htm>
2. Williamson WE, Demmler GJ, Percy AK, Catlin FI. Progressive hearing loss in Infants with asymptomatic congenital cytomegalovirus infection. *Pediatrics*. 1992;90:862-866.
3. Rivera, L.B. et al., Predictors of hearing loss in children with symptomatic congenital Cytomegalovirus infection. *Pediatrics*, 2002, 110, 762-767.
4. March of Dimes, Rubella Fact Sheet, http://www.marchofdimes.com/professionals/14332_1225.asp
5. March of Dimes, Sexually Transmitted Infections in Pregnancy Fact Sheet, http://www.marchofdimes.com/professionals/14332_1226.asp
6. March of Dimes, Toxoplasmosis Fact Sheet, http://www.marchofdimes.com/professionals/14332_1228.asp
7. March of Dimes, Sexually Transmitted Infections in Pregnancy Fact Sheet, http://www.marchofdimes.com/professionals/14332_1226.asp
8. Ely, JW et al., Evaluation of Pregnant Women Exposed to Respiratory Viruses. *American Family Physician*, 2000; 61, 10. <http://www.aafp.org/afp/20000515/3065.html>
9. Kountakis, SE. Risk factors for hearing loss in neonates: a prospective study. *American Journal of Otolaryngology*. 2002 May-Jun;23(3):133-7.
10. Greater Baltimore Medical Center Healthcare <http://www.gbmc.org/genetics/harveygenetics/prenataldx/PatientInformation/teratogens.cfm>

For more information about infant hearing loss, please visit our website:
www.doh.wa.gov/EarlyHearingLoss/Provider

Contact us at:

Washington State Department of Health
Early Hearing Loss, Detection, Diagnosis, and Intervention (EHDDI) Program
1610 NE 150th Street, K17-9, Shoreline, WA 98155-0729
Phone: (206) 418-5613 / 1-888-WAEHDDI (1-888-923-4334)
E-mail: ehddi2@doh.wa.gov

